

The Fate of Sulfur and Halogen Gases in a Plinian Volcanic Plume

PETER M. SACHS, GEOMAR, VOLCANOLOGY AND PETROLOGY, KIEL CHRISTIANE TEXTOR, HANS-F. GRAF, MAX-PLANCK-INSTITUTE FOR METEOROLOGY, HAMBURG



MOTIVATION

THE MASS OF VOLCANIC HALOGENS INJECTED INTO THE STRATOSPHERE IS UNKNOWN AT PRESENT!

Volcanic emissions in the stratosphere: effect the climate for up to three years,

destroy stratospheric ozone
 Pinatubo (1991) over Europe, North America and Central Africa: 10-20%.

Sulfur gases (SO2, H2S)

aerosols strongly increase ozone-depleting potential of halogen species


OBJECTIVES

- To what extent do volcanic gases from a typical plinian eruption reach the stratosphere? Which processes in the eruption column determine the scavenging efficiency?

METHOD

Numerical simulation of the Plinian eruption of the Laacher See volcano, Germany, 12900 BP, with the plume model ATHAM.

Laacher See volcano





Estimate of volatile release of F, CI, Br, I, SO₂, and H₂O

Chlorine, Rustine, sulfur.
The eruptive released of the 40 SQ can not be calculated by application of the presence of the calculated by application of the period (Devine et al., 1994), due to a late magma contamination (prograde fertilization and amphibole and probably sulfide break-down) (Sachs and Harms, 2010).

The clearest evidence for the syneruptive atmospheric emission is pro-vided by bromine. The mass of Bris determined by the petrologic method.

Mass of released F, Cl, I, SO₂ and H₂O: $M = M_{Br}(D_x/D_{Br})(C_x/C_{Br})$.

 $\begin{array}{lll} C_{\rm D}/C_{\rm Br} &= 64..85 \\ C_{\rm F}/C_{\rm gr} &= 41..55 \\ C_{\rm J}/C_{\rm gr} &< 0.003 \\ C_{\rm SQ}/C_{\rm gr} &< 18..25 \\ C_{\rm H2O}/C_{\rm Br} &= 65500/35 = 1871 \\ \end{array}$

If inpited mass of Br or components x which can be F, Cl.1, SO_2 or H_2O . C Preequiptive equilibrium concentrations in the meti, represented at best through gioss inclusions which have the highest concentrations of $to (C_{to}-3, d)_{to}(g/s)$. D Position coefficients between fluid and met (Wester 1990, Keppler 1999, Bureau et al., $200, D_{to}Q_0 = 1000, S_0 = 102$) thems and Gardner, posis, comm.)

component	new method	petrologic method
Br	0.380.39	0.380.39
CI	11.515.1	6.6
1	<0.007	-
F	0.160.22	<0
SO ₂	18.926.2	3.8
H ₂ O	743558	403

The ratio of the mass of SO $_2$ relative to the volume of the exupted magma (19.9-27.6 Tg SO $_2$, 5.6 km² DRE magma) is similar to the emissions of the Phathub eruption (\sim 1 T $_2$ SO $_2$ in the startosphere (TOMS), 3.7-5.3 km² DRE magma, a cifed in Kress (1977)).

GAS SCAVENGING

Solution of gases in water drops

Kinetics of the phase transfer+ dissociation of acidic gases in water (Schwartz 1986)





 $\frac{\Delta c_{aq,i}}{\Delta t} = k_{\uparrow} L(c_{g,i} - c_{g,i}, eq)$ caqJ conc. of species i in water conc. of species i in gas cg.J.eq equilibrium gas conc. phase transfer coefficient liquid water content

Incorporation of volcanic gases into ice particles

Transfer of volcanic gases proportional to water vapour transfer (Dominé et d. 1995, Dominé and Thibert 1996) Concentration of gases in ice is determined by the kinetics of condensation





Transfer of solutes during microphysical processes

Time dependent change of chemical species in hydrometeors: proportional to the change of hydrometeors caused by different

$$\frac{\Delta c_{X}}{\Delta t} = \sum_{\text{DFOC}} \frac{\Delta q_{X}}{\Delta t} \frac{c_{X}}{q_{X}}$$

Sensitivity studies in a box model showed that photochemistry is suppressed in the eruption column due to the lack of light! HBr >Br₂ like suggested by Bureau et al. 2000, is negligible. QR Sander, pats comm)

ATHAM

Active Tracer High Resolution Atmospheric Model

Modules

Microphysics

Advection and thermodynamics Solution of the Navier-Stokes equation (Oberhuber et al. 1998, Herzog 1998) Entrainment of ambient air (Herzog 1998, Oberhuber et al. 1998)

Complete cloud microphysics Liquid water and ice Feedback on the dynamics included (Graf et al. 1999, Herzog et al. 1998, Textor 2000)

Scavenging of Particles by hydrometeors Particle growth due to interaction with hydrometeors (Textor et al. 2000)

Simulation Conditions

he eruption conditions correspond to the Laacher See volcano

Model version

findrical coordinates

126 x 125 grid points 150 x 50 km², focusing grid (Δx_{min}=50m)

Geometry of the volcano mountain height (asl) diameter of the crater depth of the crater

Volcanic forcina

Composition of the magma 95 normalised gamma distribution 2 classes 40 % radius=25 µm 60 % radius=50 µm H₂O SO₂ HCI HBr

Atmospheric profiles for temperature and relative humidity Midlatitudes



Simulation Results for Hydrometeors

-45-40-35-30-25-39-15-10-5 0 5 10 15 29 25 30 35 40 Distance [km] Total condensed water and ice mass mixing ratio (g/kg) Frozen hydrometeors are highly dominant in the plume
 Liquid water exists only in the central rising zone (<1%).

Conclusions

The scavenging efficiency depends on:

ndividual solubility of the species, HCI / HBr: up to 40% contained in ice

nigh solubility in liquid water accorporation in ice

SO₂ / H₂S: up to 10% contained in ice low solubility in liquid water incorporation in ice

the **amount of hydrometeors** in the plume In a wet environment the entrainment of humid ambient air increases the hydrometeor content. This might strongly enhance the scavenging of volcanic gases.

the amount of liquid water

A caignimbrite eruption with a slower vertical velocity might show a much higher content of liquid water, which could efficiently remove HCI from the atmosphere.

The **final amount of volcanic gases in the stratosphere** depends on the fate of the contaminated hydrometeors:

Precipitation would remove species from the stratosphere Sublimation releases species into the gas phase.


n our study, sublimation of ice releases the species back to the gas shase in the stratosphere at the end of the simulation time.


References

DEPENCION STATEMENT OF LIFE 13 THE JUST AND STATEMENT OF LIFE 13 THE JUST

Volcanic Particles

30 min. after eruption end







- The neutral buoyancy height around 15 km, well in the stratosphere
 Sedimentation removes particles from the stratosphere.


 -30 % or the particles are deposited 30 min, after the eruption end.


total HCl. 30 min. of eruption total HCI, 30 min, after eruption end

- The neutral buoyancy height around 15 km in the stratosphere
 Most of the HCL stays at the neutral buoyancy height after the
 Gas particle separation is very efficient.











- HCl is contained in all phases of hydrometeors.
 Despite its high water solubility the scavenging by liquid water is not very significant.

Distribution of volcanic species in different phases



- More than 99 % of the total erupted HCl and SO₂ reach the stratosphere
- \bullet The species are partly contained in ice (HCI up to 40 %, SO $_{\!2}$ up to 10 %). \bullet Precipitation of graupel from the stratosphere is insignificant (particle size $\sim 1 \mu m$).

- freezing of these drops transfers dissolved HCl to ice particles,
- HCl is incorporated in growing ice,
- is not dissolved in liquid drops.
- is incorporated in growing ice
- the sublimation totally releases $5O_2$ back



We wish to thank Rolf Sander, Max-Planck-Institute for Chemistry, Mainz, for his helpful cooperation.